

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 86103407.2

(22) Date of filing: 13.03.86

(51) Int. Cl.⁴: **C 07 D 241/12**
C 07 D 409/06, A 61 K 31/4-95

(30) Priority: 15.03.85 JP 52115/85

(43) Date of publication of application:
17.09.86 Bulletin 86/38

(84) Designated Contracting States:
BE CH DE FR GB IT LI NL SE

(71) Applicant: **TERUMO KABUSHIKI KAISHA** trading as
TERUMO CORPORATION
44-1, 2-chome, Hatagaya Shibuya-Ku
Tokyo 151(JP)

(72) Inventor: **Wakabayashi, Toshio**
30-30-2, 5-chome, Nagayama
Tama-shi Tokyo(JP)

(72) Inventor: **Hasegawa, Hirokazu**
14-4, 8-chome, Jindaijihigashimachi
Chofu-shi Tokyo(JP)

(72) Inventor: **Ohta, Akihiro**
10-18, 3-chome, Nishikolwa
Edogawa-ku Tokyo(JP)

(74) Representative: **Henkel, Feller, Hänzel & Partner**
Möhlstrasse 37
D-8000 München 80(DE)

(54) **Pyrazine derivatives.**

(57) Novel pyrazine derivatives are disclosed. The compounds possess a potent platelet aggregation-inhibiting activity and are effective for preventing diseases caused by aggregation of the platelet, for example, myocardial infarction and cerebral thrombosis. Representative examples of the pyrazine derivatives include 2,3-bis (p-chlorophenyl)-5-methylpyrazine, 2,3-bis(p-methoxyphenyl)-5-methylpyrazine, 2,3-diphenyl-5-benzylpyrazine, 2,3-diphenyl-5-(p-methoxybenzyl) pyrazine, 2,3-bis (p-methoxyphenyl)-5-isopropylpyrazine, 2,3-bis(p-methoxyphenyl)-5-(2-thienylmethyl) pyrazine and 2,3-bis-(p-methoxyphenyl)-5,6-dimethylpyrazine.

The pyrazine derivatives can be generally prepared by heating a benzil derivative and a 1, 2-diaminoethane derivative to produce a dihydropyrazine derivative and heating the resulting compound with sulfur at 100 - 180°C.

EP 0 194 686 A1


Title of the Invention

Pyrazine derivatives

Background of the Invention

1. Field of the Invention

5 This invention relates to novel pyrazine derivatives.
More particularly, it is concerned with pyrazine
derivatives having the formula



(wherein X represents a hydrogen atom, a halogen atom,
a lower alkyl group, a lower alkoxy group or a di-lower alkyl-
10 amino group, R¹ represents a hydrogen atom or a lower
alkyl group and R² represents a lower alkyl group, a
benzyl group, a substituted benzyl group having as the
substituent a lower alkyl group, a lower alkoxy group or
a methylenedioxy group, a thienylmethyl group or a
15 substituted thienylmethyl group having as the substituent
a lower alkyl group, a lower alkoxy group or a methylene-
dioxy group.

The pyrazine derivatives (I) possess a potent
platelet aggregation-inhibiting activity. Therefore,
20 they are effective for preventing diseases caused by

aggregation of the platelets, that is, such diseases as myocardial infarction and thrombosis. The pyrazine derivatives (I) of the invention also have a cyclooxygenase-inhibiting activity. As compounds with such activity
5 are generally known to possess an antiinflammatory activity, the pyrazine derivatives (I) are expected to find use as the antiinflammatory agent.

2. Description of the Prior Art

There have been known various compounds which have
10 platelet aggregation-inhibiting activities. Their activities, however, are so weak that development of drugs possessing improved effects has been desired. There is also strong need for antithrombocytic agents which will effectively prevent thrombosis such as myocardial infarction and cerebral thrombosis, which recently occupy the
15 major rate of adult diseases.

Heretofore, a variety of pyrazine derivatives are known, such as, for example, 2,3-diphenylpyrazine described in Journal of Heterocyclic Chemistry, vol.
20 21, pp. 103 - 106. However, none of these pyrazine derivatives are known to possess a platelet aggregation-inhibiting activity.

Summary of the Invention

As a result of extensive studies on pharmacological activities of a variety of novel pyrazine derivatives prepared by us, we have found that specific pyrazine
5 compounds possess a potent platelet aggregation-inhibiting activity and completed the present invention.

It is therefore an object of the invention to provide novel pyrazine derivatives which are useful as an antithrombocytic agent.

10 A further object of the invention is to provide antithrombocytic agent containing as the active ingredient such pyrazine derivatives.

Detailed Description of the Invention

In accordance with the present invention there are
15 provided novel pyrazine derivatives having the above-mentioned formula (I). In the above-mentioned formula (I), X represents a hydrogen atom, a halogen atom, for example, chlorine, bromine or fluorine atom, a lower alkyl group, for example, methyl, ethyl, propyl, iso-
20 propyl, butyl or isobutyl, a lower alkoxy group, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy or isobutoxy or a di-lower alkylamino group, for example, dimethylamino, diethylamino, methylethylamino, methylpropylamino, ethylpropylamino or dipropylamino; R¹


represents a hydrogen atom or a lower alkyl group, for example, methyl, ethyl, propyl, isopropyl or butyl; and R^2 represents a lower alkyl group, for example, methyl, ethyl, propyl, isopropyl, butyl or isobutyl, a benzyl group, a substituted benzyl group having as the substituent a lower alkyl group, a lower alkoxy group or a methylenedioxy group, for example, 3- or 4-methylbenzyl, 3- or 4-methoxybenzyl, 3- or 4-ethoxybenzyl, 3- or 4-propoxybenzyl or 3,4-methylenedioxybenzyl, a thienylmethyl group, for example, 2- or 3-thienylmethyl or a substituted thienylmethyl group having as the substituent a lower alkyl group, a lower alkoxy group or a methylenedioxy group, for example, 4- or 5-methyl-2- or 3-thienylmethyl.

15 As preferred examples of the pyrazine derivatives having the above-mentioned formula (I) are mentioned:

2,3-Diphenyl-5-methylpyrazine,
2,3-bis(p-chlorophenyl)-5-methylpyrazine,
2,3-bis(p-bromophenyl)-5-methylpyrazine,
20 2,3-bis(p-methylphenyl)-5-methylpyrazine,
2,3-bis(p-methoxyphenyl)-5-methylpyrazine,
2,3-bis(p-dimethylaminophenyl)-5-methylpyrazine,
2,3-bis(p-chlorophenyl)-5-ethylpyrazine,
2,3-diphenyl-5-benzylpyrazine,
25 2,3-diphenyl-5-(p-methylbenzyl)pyrazine,

- 2,3-diphenyl-5-(p-methoxybenzyl)pyrazine,
 2,3-diphenyl-5-(m-methoxybenzyl)pyrazine,
 2,3-bis(p-methoxyphenyl)-5-benzylpyrazine,
 2,3-bis(p-chlorophenyl)-5-benzylpyrazine,
 5 2,3-bis(p-methoxyphenyl)-5-isopropylpyrazine,
 2,3-bis(p-methoxyphenyl)-5-ethylpyrazine,
 2,3-bis(p-methoxyphenyl)-5-(2-thienylmethyl)pyrazine,
 2,3-diphenyl-5-(2-thienylmethyl)pyrazine,
 2,3-bis(p-chlorophenyl)-5-(2-thienylmethyl)pyrazine and
 10 2,3-bis(p-chlorophenyl)-5-(4- or 5-methyl-2- or 3-thienyl-
 methyl)pyrazine.


The pyrazine derivatives (I) are prepared by heating
 a benzil derivative having the formula



- wherein X has the same meaning as described above and
 15 a 1,2-diaminoethane derivative having the formula



wherein R^1 and R^2 have the same meaning as described above in an appropriate organic solvent such as, for example, ethanol to produce a dihydropyrazine derivative having the formula



5 wherein X, R^1 and R^2 have the same meaning as described above and subsequently heating the same with sulfur at 100 - 180°C.

The pyrazine derivatives (I) wherein R^1 is a hydrogen atom and R^2 is a benzyl, substituted benzyl, thienyl-
 10 methyl or substituted benzyl group may also be prepared by reacting the above-described benzil derivatives with 1,2-diaminoethane to produce a dihydropyrazine derivative and reacting the latter compound with a substituted or unsubstituted benzaldehyde or thiophenealdehyde. In
 15 the above preparative process, when a dialkyl ketone is reacted in place of the benzaldehyde the pyrazine derivatives (I) wherein R^2 is a branched chain-alkyl group can be produced. Furthermore, pyrazine derivatives of the above-described formula (I) wherein R^1 and R^2 are
 20 a lower alkyl group, respectively may also be obtained by oxidizing a 2,3-diphenylpyrazine derivative with

permaleic acid to produce a mono- or di-N-oxide product, then chlorinating the same with phosphorus oxychloride to convert to a 2,3-diphenyl-4-(or 4,5)-chloropyrazine and reacting the same with a trialkylboron.

5 As the pyrazine derivatives (I) of the invention possess a platelet aggregation-inhibiting activity, they are effectively utilized as an antithrombocytic agent for prevention of cerebral thrombosis and like diseases. Moreover, the pyrazine derivatives (I) of the invention
10 possess a cyclooxygenase-inhibiting activity and are utilizable as an antiinflammatory agent.

 The pyrazine derivatives of the invention may be administered at a dosage level in a range between about 30 and 600 mg per day in adults, if necessary, divided
15 into one to three doses. The route of administration may be in any form suitable for administration, oral administration being particularly desirable with intravenous administration also feasible.

 The compounds of the invention are formulated
20 either alone or in admixture with pharmaceutical carriers or excipients by a conventional method into tablet, powder, capsule or granule. As examples of the carrier or excipient are mentioned calcium carbonate, starch, sucrose, lactose, talc, magnesium stearate and the like.
25 In addition to the above-mentioned solid preparations,

the compounds of the invention may also be formulated into liquid preparations such as oily suspension or syrup. They may also be stabilized in the form of inclusion in cyclodextrin.

5 The following examples and test examples are intended to illustrate the invention more concretely, but are not to be construed as limiting the scope thereof.

Example 1.

10 To a solution of 2.09 g of 4,4'-di-chlorobenzyl in 35 ml of ethanol was added dropwise 0.66 g of 1,2-propanediamine at room temperature. The mixture was heated under reflux for 30 minutes. The reaction mixture was concentrated to half of its original volume under reduced pressure. After ice-cooling, the produced
15 precipitates were collected by filtration and recrystallized from ethanol to give 1.53 g of 2,3-bis(p-chlorophenyl)-5,6-dihydro-5-methylpyrazine as yellow needles with a melting point of 125 - 126°C. 930 mg of the compound was mixed with 192 mg of powdery sulfur and
20 the mixture was heated on an oil bath at 140°C for 30 minutes. The reaction mixture was allowed to cool and subjected to column chromatography on silica gel. 630 mg of 2,3-bis(p-chlorophenyl)-5-methylpyrazine was obtained from the eluates with benzene. Physicochemical

properties of the product support a chemical structure of the below formula (V).

M.P. 168 - 169°C (recrystallized from n-hexane).


Analysis:

5 Calc'd. for $C_{17}H_{12}N_2Cl_2$: C, 64.78%; H, 3.84%;
N, 8.89%.

Found: C, 64.42%; H, 3.82%; N, 8.78%.

Mass (m/e): 314 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 2.60 (3H, s), 8.40 (1H, s).



(V)

10 Example 2

To a solution of 5.40 g of 4,4'-dimethoxybenzyl in 100 ml of ethanol was added dropwise 1.78 g of 1,2-propanediamine at room temperature. The mixture was heated under reflux for an hour. The reaction mixture
15 was allowed to cool and insolubles precipitated were filtered off. The solvent was distilled off under reduced pressure from the mother liquor. The resulting residue was subjected to column chromatography on silica gel to obtain 3.549 g of 2,3-bis(p-methoxyphenyl)-5,6-
20 dihydro-5-methylpyrazine from the eluate with a 1 : 1

n-hexane : methylene chloride mixture. To a solution of 1.004 g of the resulting product in 20 ml of methylene chloride was added 208 mg of powdery sulfur. The methylene chloride was distilled off under reduced pressure from the reaction mixture and the resulting residue was heated on an oil bath at 140°C for 15 minutes. The reaction mixture was allowed to cool and subjected to column chromatography on silica gel to give 754 mg of 2,3-bis(p-methoxyphenyl)-5-methylpyrazine from the eluates with benzene. Physicochemical properties of the product support a chemical structure of the below formula (VI).

M.P. 120 - 121°C (recrystallized from methanol).


Analysis:

Calc'd. for $C_{19}H_{18}N_2O_2$: C, 74.49%; H, 5.92%; N, 9.14%.

Found: C, 74.63%; H, 6.20%; N, 9.12%.

Mass (m/e): 306 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 2.60(3H,s), 3.77(3H,s), 6.77(2H,dd,J=2Hz,10Hz), 7.42(2H,dd,J=2Hz,10Hz), 8.40(1H,s).



(VI)

Example 3

In 20 ml of methanol were dissolved 2.570 g of 2,3-diphenyl-5,6-dihydropyrazine, 1.060 g of benzaldehyde and 0.672 g of potassium hydroxide. The solution was heated under reflux for 1 hour. Methanol was removed from the reaction solution by distillation under reduced pressure, and to the residue was added 50 ml of water. The resulting mixture was extracted three times with ethyl acetate. The organic layer from the extraction was washed with water and dried over anhydrous sodium sulfate. The solvent was removed by distillation under reduced pressure. The residue thus obtained was subjected to column chromatography on silica gel. A crude product was obtained from the eluates with an 8 : 1 hexane : ethyl acetate mixture. The crude product was recrystallized from hexane to give 2.940 g of 2,3-diphenyl-5-benzylpyrazine as colorless prisms. M.P. 118 - 119°C. Physical properties of the product support a chemical structure of the below formula (VII).


Analysis:

Calc'd. for $C_{23}H_{18}N_2$: C, 85.68%; H, 5.63%;
N, 8.69%.

Found: C, 85.68%; H, 5.53%; N, 8.67%.

Mass (m/e): 322 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 4.27(2H,s), 8.47(1H,s).



(VII)

Example 4

The same procedures as in Example 3 were repeated using 2,3-diphenyl-5,6-dihydropyrazine and p-anisaldehyde. 2,3-Diphenyl-5-(p-methoxybenzyl)pyrazine was obtained as colorless prisms, m.p. 102 - 103°C (recrystallized from methanol-water). Physical properties of the product support a chemical structure of the below formula (VIII).


Analysis:

Calc'd. for $C_{24}H_{20}N_2O$: C, 81.79%; H, 5.72%;
N, 7.95%.

Found: C, 81.89%; H, 5.70%; N, 8.01%.

Mass (m/e): 352 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 3.73(3H,s), 4.17(2H,s),
6.83(2H,d,J=9Hz), 8.37
(1H,s).



(VIII)

Example 5

The same procedures as in Example 3 were repeated using 2,3-diphenyl-5,6-dihydropyrazine and m-anisaldehyde. 2,3-Diphenyl-5-(m-methoxybenzyl)pyrazine was obtained as colorless prisms, m.p. 65 - 66°C (recrystallized from methanol-water). Physical properties of the product support a chemical structure of the below formula (IX).


Analysis:

Calc'd. for $C_{24}H_{20}N_2O$: C, 81.79%; H, 5.72%;
N, 7.95%.

Found: C, 82.02%; H, 5.74%; N, 7.97%.

Mass (m/e): 352 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 3.73(3H,s), 4.18(2H,s),
8.38(1H,s).



Example 6

The same procedures as in Example 3 were repeated using 2,3-bis(p-methoxyphenyl)-5,6-dihydropyrazine and benzaldehyde. 2,3-Bis(p-methoxyphenyl)-5-benzylpyrazine was obtained as colorless prisms, m.p. 107-109°C (recrystallized from methanol-water). Physical properties of the product support a chemical structure of the below

formula (X).


Analysis:

Calc'd. for $C_{25}H_{22}N_2O_2$: C, 78.51%; H, 5.80%;
N, 7.33%.

5 Found: C, 78.48%; H, 5.77%; N, 7.29%.

Mass (m/e): 382 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 3.77(6H,s), 4.22(2H,s),
6.80(2H,d,J=9Hz), 6.82
(2H,d,J=9Hz), 8.33(1H,s).



10 Example 7

The same procedures as in Example 3 were repeated using 2,3-bis(p-methoxyphenyl)-5,6-dihydropyrazine and acetone. 2,3-Bis(p-methoxyphenyl)-5-isopropylpyrazine as an oily substance, b.p. 211°C (0.15 Torr). Physical
15 properties of the product support a chemical structure of the below formula (XI).


Analysis:

Calc'd. for $C_{21}H_{22}N_2O_2$: C, 75.42%; H, 6.63%;
N, 8.38%.

20 Found: C, 76.19%; H, 6.76%; N, 8.48%.

Mass (m/e): 334 (molecular ion peak).

$^1\text{H-NMR}$ (CDCl_3) δ (ppm): 1.37(6H,d,J=7Hz), 3.75(6H,s),
6.78(4H,d,J=9Hz), 7.35(2H,
d,J=9Hz), 7.41(2H,d,J=9Hz),
8.37(1H,s).



(XI)

5 Example 8

To 20 ml of phosphorus oxychloride was added with stirring 3.918 g of 2,3-bis(p-methoxyphenyl)pyrazine oxide. The mixture was heated under reflux for 30 minutes. After allowed to cool, the reaction mixture
10 was poured into ice water and then made basic with potassium carbonate to precipitate the product, which was collected by filtration. There was obtained 2.823 g of 2,3-bis(p-methoxyphenyl)-5-chloropyrazine. A mixture of 653 mg of said compound, 414 mg of anhydrous potassium
15 carbonate and 116 mg of tetrakis(triphenylphosphine)-palladium was suspended in an atmosphere of argon in 10 ml of dry N,N-dimethylformamide. To the suspension was added 2 ml of a hexane solution of triethylborane (15%). The resulting mixture was heated under reflux
20 for 2 hours, followed by removal of the solvent by distillation under reduced pressure. Water was added

to the residue, and the resulting mixture was extracted three times with methylene chloride. The organic layer from the extraction was washed with water and dried over anhydrous sodium sulfate. Removal of the solvent by
5 distillation under reduced pressure afforded 850 mg of an extraction residue. The residue was subjected to column chromatography on silica gel. There was obtained 555 mg of 2,3-bis(p-methoxyphenyl)-5-ethylpyrazine from the eluates with a 1 : 1 hexane : methylene chloride
10 mixture. M.P. 76.5 - 78°C (colorless needles, recrystallized from ethanol). Physical properties of the product support a chemical structure of the below formula (XII).


Analysis:

15 Calc'd. for $C_{20}H_{20}N_2O_2$: C, 74.98%; H, 6.29%;
N, 8.74%.

Found: C, 75.02%; H, 6.41%; N, 8.73%.

Mass (m/e): 320 (molecular ion peak).

20 1H -NMR ($CDCl_3$) δ (ppm): 1.37(3H,t,J=7Hz), 2.87(2H,
q,J=7Hz), 3.77(6H,s), 6.77
(4H,d,J=7.5Hz), 7.33(2H,d,
J=7.5Hz), 7.37(2H,d,J=7.5Hz),
8.33(1H,s).



(XII)

Example 9

The same procedures as in Example 3 were repeated using 2,3-bis(p-methoxyphenyl)-5,6-dihydropyrazine and 2-thiophenealdehyde. 2,3-Bis(p-methoxyphenyl)-5-(2-thienylmethyl)pyrazine was obtained as colorless prisms, m.p. 89°C (recrystallized from methanol). Physical properties of the product support a chemical structure of the below formula (XIII).


Analysis:

Calc'd. for C₂₃H₂₀N₂O₂S: C, 71.11%; H, 5.19%;
N, 7.21%.

Found: C, 71.19%; H, 5.25%; N, 7.22%.

Mass (m/e): 388 (molecular ion peak).

¹H-NMR (CDCl₃) δ (ppm): 3.77(6H,s), 4.40(2H,s),
6.80(2H,d,J=9Hz); 6.92(2H,
d,J=9Hz), 7.37(2H,d,J=9Hz),
7.43(2H,d,J=9Hz), 8.40(1H,s).



Example 10

To 20 ml of phosphorus oxychloride was added with stirring 1,800 g of 2,3-bis(p-methoxyphenyl)pyrazine 1,4-dioxide. The mixture was heated under reflux for 5 1 hour. After allowed to cool, the reaction mixture was poured into ice water and made basic with potassium carbonate. The resulting mixture was extracted three times with methylene chloride. The organic layer from the extract was washed with water and dried over anhy- 10 drous sodium sulfate. The solvent was then removed by distillation under reduced pressure to obtain an extrac- tion residue. The residue was subjected to column chromatography on silica gel to give 1.504 g of 2,3- bis(p-methoxyphenyl)-5,6-dichloropyrazine.

15 In 10 ml of dry tetrahydrofuran were dissolved in an atmosphere of argon 903 mg of said compound and 289 mg of tetrakis(triphenylphosphine)palladium. To the solution was further added 1.7 ml of a hexane solution of trimethylaluminum (15%). Then, the resulting mixture 20 was heated under reflux for 4 hours. After allowed to

cool, 1 ml of water was added to the reaction mixture,
 followed by removal of the solvent by distillation under
 reduced pressure. To the residue was again added 1 ml
 of water, and the mixture was extracted three times with
 5 methylene chloride. The organic layer from the extrac-
 tion was washed with water and dried over anhydrous
 sodium sulfate. When the solvent was removed by distil-
 lation under reduced pressure, there was produced 1.30 g
 of an extraction residue. The residue was subjected to
 10 column chromatography on silica gel. There was obtained
 694 mg of 2,3-bis(p-methoxyphenyl)-5,6-dimethylpyrazine
 from the eluates with methylene chloride. M.P. 106.5 -
 108°C (colorless needles, recrystallized from ethanol).
 Physical properties of the product support a chemical
 15 structure of the below formula (XIV).


Analysis:

Calc'd. for $C_{20}H_{20}N_2O_2$: C, 74.98%; H, 6.29%;
 N, 8.74%.

Found: C, 74.89%; H, 6.25%; N, 8.90%.

20 Mass (m/e): 320 (molecular ion peak).

1H -NMR ($CDCl_3$) δ (ppm): 2.57(6H,s), 3.77(6H,s),
 6.78(4H,d,J=7.5Hz), 7.37
 (4H,d,J=7.5Hz).



(XIV)

Pharmacological Test Example 1

Platelet Aggregation-Inhibiting Action


Nine volumes of blood was drawn from the carotid artery of a rabbit using a syringe containing one volume of 3.8% solution of sodium citrate. The blood was centrifuged to obtain platelet-rich plasma (PRP: 7×10^5 platelets/ μ l).

In a cuvette was placed 268 μ l of the PRP, which was warmed in an isothermic bath at 37°C for 2 minutes. To the cuvette was added 2 μ l of an ethanol solution of a pyrazine derivative to be tested, followed by an incubation for 3 minutes. To the incubate was added a solution of a platelet-aggregation inducer, arachidonic acid or collagen. Measurement was made by Born's turbidimetric method (for example, see J. Physiol., vol. 168, p. 178, 1968). 50% inhibitory concentration for the platelet aggregation caused by arachidonic acid (50 μ mol.) or collagen (10 μ g/ml) was given in Table 1 using acetylsalicylic acid as the reference.

As shown in Table 1, the pyrazine derivatives of

the invention were found to possess a marked platelet aggregation-inhibiting activity. The pyrazine derivatives not shown in Table 1 was also confirmed to possess a similar platelet aggregation-inhibiting activity. The
5 50% inhibitory concentration as shown in table means the concentration of a solution of the pyrazine derivative required for inhibiting the platelet aggregation to 50% when the platelet aggregation in the absence of a pyrazine derivative is taken as 100%.

Table 1. Platelet Aggregation Inhibiting Activity



Example No.	Substituent			50% aggregation inhibitory concentration (mol.)	
	X	R ¹	R ²	Arachidonic acid	Collagen
1	-Cl	H	-CH ₃	8.5 x 10 ⁻⁶	2.0 x 10 ⁻⁵
2	-OCH ₃	H	-CH ₃	3.0 x 10 ⁻⁸	4.2 x 10 ⁻⁸
3	H	H	-CH ₂ -	8.0 x 10 ⁻⁶	4.6 x 10 ⁻⁵
4	H	H	-CH ₂ -	1.7 x 10 ⁻⁶	3.5 x 10 ⁻⁵
5	H	H	-CH ₂ -	4.5 x 10 ⁻⁶	1.4 x 10 ⁻⁵
6	-OCH ₃	H	-CH ₂ -	2.0 x 10 ⁻⁷	5.0 x 10 ⁻⁶
7	-OCH ₃	H	-CH(CH ₃) ₂	2.5 x 10 ⁻⁸	7.9 x 10 ⁻⁷
8	-OCH ₃	H	-C ₂ H ₅	4.6 x 10 ⁻⁸	6.4 x 10 ⁻⁷
9	-OCH ₃	H	-CH ₂ -	4.4 x 10 ⁻⁷	7.9 x 10 ⁻⁷
10	-OCH ₃	-CH ₃	-CH ₃	8.4 x 10 ⁻⁸	9.2 x 10 ⁻⁷
Aspirin (Control)	-	-	-	1.4 x 10 ⁻⁵	5.6 x 10 ⁻⁴

Test Example 2

Cyclooxygenase Inhibiting Activity

Nine volumes of blood was drawn from the abdominal artery of a rabbit using a syringe containing one volume of 3.8% solution of sodium citrate. Centrifugation of the blood afforded platelet-rich plasma. To the platelet-rich plasma was added 77 mM EDTA solution in a volume of 1/10 per volume of the plasma. The mixture was thoroughly mixed and centrifuged at 2500 rpm for 10 minutes. The supernatant was discarded, and the platelets were suspended in approximately 3 ml of a washing solution which was prepared by dissolving 134 mM of sodium chloride, 15 mM of trisaminomethane, 1 mM of EDTA and 5 mM of D-glucose in twice-distilled water with a pH adjusted with 1N hydrogen chloride to 7.4. The suspension was centrifuged at room temperature at 2000 rpm for 10 minutes. The supernatant was discarded, and the precipitated platelets were re-suspended in a phosphate buffer at pH 8.0 to adjust number of the platelets to 1×10^6 per μl .

The washed platelets thus obtained were employed as a cyclooxygenase source.

To 3 μg of arachidonic acid and 0.2 μCi (1 μg) of ^{14}C -labelled arachidonic acid placed in a glass-stoppered test tube was added one drop of a propylene



glycol/ethanol solution (1 : 3 by volume). The ethanol was evaporated under nitrogen. To the residue were added 50 μ l of a solution to be tested and then 450 μ l of the washed platelets. The mixture was reacted at
5 37°C for 3 minutes.

To the reaction mixture was added with ice cooling one drop of 1N hydrogen chloride to adjust the pH to 2 - 3. To the resulting mixture was added 2 ml of ethyl acetate, followed by extraction by shaking for 10 minutes
10 and centrifugal separation at 4°C at 2500 rpm for 10 minutes.

The supernatant was transferred to a flask and concentrated. The residue was dissolved in 100 μ l of ethanol, and the entire solution was spotted on a silica
15 gel thin plate (Merck, 60 F₂₅₄).

The plate was developed by approximately 18 cm with a developer solution (chloroform/methanol/acetic acid/waters = 70 : 8 : 1 : 0.8), followed by measurement by a radiochromatoscanner of the sum of radioactivities
20 of prostaglandin F_{2 α} , thromboxan B₂, prostaglandin E_{2 α} , prostaglandin D₂ and HHT to determine the inhibiting activity. Results are shown in Table 2.

Table 2. Cyclooxygenase Inhibiting Activity


Example No.	Substituent			50% Inhibitory concentration (mol.)
	X	R ¹	R ²	
2	-OCH ₃	H	-CH ₃	3.0 x 10 ⁻⁵
4	H	H	-CH ₂ - 	2.2 x 10 ⁻⁴
9	-OCH ₃	H	-CH ₂ - 	8.6 x 10 ⁻⁵
10	-OCH ₃	-CH ₃	-CH ₃	6.3 x 10 ⁻⁵

Acute Toxicity

An acute toxicity test was conducted in male ICR mice (5 week old) by oral administration. LD₅₀ values were 300 mg/Kg or higher for all of the pyrazine derivatives of the invention tested thereby demonstrating high safety.

What is claimed is:

1. A pyrazine derivative having the formula



wherein X represents a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group or a di-loweralkyl-amino group, R^1 represents a hydrogen atom or a lower alkyl group and R^2 represents a lower alkyl group, a benzyl group, a substituted benzyl group having as the substituent a lower alkyl group, a lower alkoxy group, a methylenedioxy group or a thienylmethyl group or a substituted thienylmethyl group having as the substituent a lower alkyl group, a lower alkoxy group or a methylene-dioxy group.

2. The pyrazine derivative according to Claim 1 wherein X represents a hydrogen atom, a halogen atom, a lower alkyl group or a lower alkoxy group, R^1 represents a hydrogen atom or a lower alkyl group and R^2 represents a lower alkyl group, a benzyl group, a substituted benzyl group having as the substituent a lower alkoxy group.

3. A platelet aggregation inhibitor containing a medicinally effective amount of an active ingredient of the pyrazine derivative according to Claim 1.
4. A platelet aggregation inhibitory pharmaceutical composition comprising a therapeutically effective amount of the pyrazine derivative according to Claim 1 and a pharmaceutical carrier therefor.
5. A method for preventing a disease caused by aggregation of the platelet which comprises administering a therapeutically effective amount of the pyrazine derivative according to Claim 1 to a mammal possibly afflicted with said disease.
6. A method of preventing a disease of Claim 5 wherein the disease is thrombosis.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

0194686
Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 86103407.2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	CHEMICAL ABSTRACTS, vol. 96, no. 9, March 1, 1982, Columbus, Ohio, USA Yamanaka, Hiroshi; Konno, Shoetsu; Sakamoto, Takao; Niitsuma, Setsuko; Noji, Sayo. "Studies on pyrimidine derivatives. XXIII. Synthesis of acylmethylpyrimidines and related compounds via imido-yl-substituted oxosulfonium ylides". * Page 607, column 2, abstract-no. 68 932t * & Chem. Pharm. Bull. 1981 29(10), 2837-43	1	C 07 D 241/12 C 07 D 409/06 A 61 K 31/495
A	CHEMICAL ABSTRACTS, vol. 83, no. 5, August 4, 1975, Columbus, Ohio, USA Padwa, Albert; Gehrlein, Lane; Kinnel, Robin B. "Synthesis of and base-induced rearrangements in the 1,4-diazabicyclo [4.1.0] hept-4-ene system".	1	
INCOMPLETE SEARCH			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims. Claims searched completely: 1-4 Claims searched incompletely: - Claims not searched: 5,6. Reason for the limitation of the search: Article 52(4) EPC; method for treatment of the human or animal body by therapy.			C 07 D 241/00 C 07 D 409/00
Place of search VIENNA		Date of completion of the search 14-05-1986	Examiner HAMMER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons a : member of the same patent family, corresponding document



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

0194686

EP 86103407.2

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	<p>* Page 498, column 1, abstract- no. 43 267y *</p> <p>& J.Org.Chem. 1975, 40(12), 1683-8</p> <p>--</p> <p>A CHEMICAL ABSTRACTS, vol. 99, no. 11, 1 September 12, 1983, Columbus, Ohio, USA</p> <p>Joshi, S.C.; Mehrotra, K.N. "A new synthesis of pyrazines".</p> <p>* Page 569, column 1, abstract- no. 88 170h *</p> <p>& Indian J.Chem., Sect. B 1983, 22B(4), 396-7</p> <p>--</p> <p>A CHEMICAL ABSTRACTS, vol. 102, no. 1 1, January 7, 1985, Columbus, Ohio, USA</p> <p>Ohta, Akihiro; Inoue, Akira; Watanabe, Tokuhiko "Introduction of the methyl group into the pyra- zine ring".</p> <p>* Page 584, column 2, abstract- no. 6 421j *</p> <p>& Heterocycles 1984, 22(10), 2317- 21</p> <p>----</p>		
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.